



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# THE AMERICAN MATHEMATICAL MONTHLY.

Entered at the Post-office at Springfield, Missouri, as second-class matter.

VOL. X.

MAY, 1903.

No. 5.

## ON THE PRODUCT OF AN ALTERNANT BY A SYMMETRIC FUNCTION.

By DR. W. E. TAYLOR, Associate Professor of Mathematics, Syracuse University.

As is well known, the product of the simple alternant  $| a_1^0 a_2^1 a_3^2 a_4^3 \dots |$  and a symmetric function of  $a_1, a_2, a_3, \dots$  is an aggregate of alternants. When the symmetric function is of the form  $\sum a_1 a_2 a_3 \dots a_r$ , the product is a single alternant differing from the original in having each of the last exponents increased by unity. In the general case, the mode of obtaining the aggregate is a fairly simple problem. The problem of obtaining the coefficient of a given alternant without having at the same time to obtain the coefficients of all the alternants in the aggregate is a problem not so simple. As a partial solution of this problem Dr. Muir has shown how the coefficient of one term of the aggregate may be obtained independently of the others.

The object of this paper is to extend the general problem started by Muir.

It is apparent that if we have a table giving the coefficients of all alternants in the product, (1)

$$| 0123 \dots (n-1) | \left( \sum a_1 a_2 a_3 \dots a_{j_1} \right)^{i_1} \left( \sum a_1 a_2 a_3 \dots a_{j_2} \right)^{i_2} \dots \left( \sum a_1 a_2 a_3 \dots a_{j_t} \right)^{i_t}$$

where  $0 \leq i_k \leq n$ ,  $1 \leq j_h \leq n$ ;  $i_1 j_1 + i_2 j_2 + \dots + i_t j_t = t \leq n$ ; and where  $i_1 + i_2 + i_3 + \dots + i_h = k$ , we can by taking the proper multiples\* of

$$(2) \left( \sum a_1 a_2 a_3 \dots a_{j_1} \right)^{i_1} \left( \sum a_1 a_2 a_3 \dots a_{j_2} \right)^{i_2} \dots \left( \sum a_1 a_2 a_3 \dots a_{j_h} \right)^{i_h}$$

form any symmetric function  $S$ , and hence from such a table get the coefficients of any alternant in the product  $| 0 \ 1 \ 2 \ 3 \dots (n-1) | S$ .

---

\*These multiples are found from the table of symmetric functions of weight  $t$ .

The difference equation for the coefficients of all alternants in the product (1) (where  $i_1=1, j_1 \leq \frac{1}{2}n$ ) is obtained and this is used to construct tables for certain cases.

1. Let  $|0\ 1\ 2\ 3 \dots r_1 s_1 \dots r_2 s_2 \dots r_g s_g \dots r_k s_k|$  denote the alternant wherein the numbers are consecutive except at the  $k$  points  $r_1 s_1, r_2 s_2, \dots, r_k s_k$ ,—that is, the numbers from 0 to  $r_1, s_1$  to  $r_2, s_2$  to  $r_3, s_3$ , and so on are consecutive; but  $r_1 s_1, r_2 s_2$ , etc., may differ by more than unity.

Let us denote the coefficient of this alternant in the product

$$|0\ 1\ 2\ 3 \dots (n-1)| \left( \sum a_1 a_2 a_3 \dots a_{j_1} \right) \left( \sum a_1 a_2 a_3 \dots a_{j_2} \right)^{i_2} \dots \left( \sum a_1 a_2 a_3 \dots a_{j_h} \right)^{i_h}$$

by

$$(3) \quad C^t \left\{ \begin{matrix} s_1 & s_2 & s_3 & \dots & s_g & \dots & s_k \\ 0 & 0 & 0 & \dots & 0 & \dots & 0 \end{matrix} \right\}$$

and the coefficient of

$$|0\ 1\ 2\ 3 \dots r_1 (s_1-1) \dots (s_1+\alpha-1)(s_1+\alpha) \dots r_2 (s_2-1) s_2 \dots (s_2+\beta-2)(s_2+\beta) \dots$$

$$\dots r_g (s_g-1) s_g \dots (s_g+\gamma-2) \dots r_k (s_k-1)| \text{ in the product}$$

$$|0\ 1\ 2\ 3 \dots (n-1)| \left( \sum a_1 a_2 a_3 \dots a_{j_1} \right) \left( \sum a_1 a_2 a_3 \dots a_{j_2} \right)^{i_2} \dots \left( \sum a_1 a_2 a_3 \dots a_{j_h} \right)^{i_h-1}$$

by

$$C^{t-j_h} \left\{ \begin{matrix} s_1 & s_2 & \dots & s_g & \dots & s_k \\ \alpha & \beta & \dots & \gamma & \dots & \kappa \end{matrix} \right\},$$

where  $\gamma$  placed below  $s_g$  denotes that  $\gamma$  consecutive numbers beginning with  $s_g$  are decreased by unity.

2. Now it is easily seen that that the product

$$|0\ 1\ 2\ 3 \dots (n-1)| \left( \sum a_1 a_2 a_3 \dots a_{j_1} \right) \left( \sum a_1 a_2 a_3 \dots a_{j_2} \right)^{i_2} \dots \left( \sum a_1 a_2 a_3 \dots a_{j_h} \right)^{i_h-1}$$

$$= \dots + \sum_0^{j_h} \alpha \beta \dots \kappa C^{t-j_h} \left\{ \begin{matrix} s_1 & s_2 & \dots & s_k \\ \alpha & \beta & \dots & \kappa \end{matrix} \right\} |0\ 1\ 2\ 3 \dots$$

$$\dots r_1 (s_1-1) s_1 (s_1+1) \dots (s_1+\alpha-2)(s_1+\alpha) \dots r_g (s_g-1) s_g \dots (s_g+\gamma-2)(s_g+\gamma) \dots$$

$$\dots r_k (s_k-1)| + \text{other terms, where } \alpha + \beta + \dots + \kappa = j_h; \kappa \text{ cannot of course be greater than 1.}$$

Only those terms are written which on multiplying both sides by  $\Sigma a_1 a_2 a_3 \dots a_{j_h}$  can give rise to

$$|0\ 1\ 2\ 3 \dots r_1 s_1 \dots r_2 s_2 \dots r_g s_g \dots r_k s_k| ;$$

but the coefficient of this term in the product is (3). Hence this coefficient must be equal to the sum of the coefficients of the above terms, that is



	$\frac{10123 \dots (n-4)(n-2)(n+1)}{10123 \dots (n-5)(n-3)(n-2)(n+1)}$ $\frac{10123 \dots (n-5)(n-3)(n-2)(n+2)}{10123 \dots (n-3)(n+3)}$ $\frac{10123 \dots (n-2)(n-9) \dots (n-1)}{10123 \dots (n-5)(n-3)(n-2)(n+3)}$ $\frac{10123 \dots (n-2)(n-1)(n+3)}{10123 \dots (n-2)(n+3)}$										
$(\sum a_i)^6$	5	9	16	9	5	1	5	10	10	5	1
$\sum a_i a_j (\sum a_i)^4$	3	6	8	3	2	1	4	6	4	1	
$(\sum a_i a_j)^4 (\sum a_i)^2$	2	4	4	1	1	1	3	3	1		
$(\sum a_i a_j)^3$	1	3	2		1	1	2	1			
$\sum a_i a_j a_k (\sum a_i)^3$	1	3	2			1	3	3	1		
$\sum a_i a_j a_k \sum a_i a_l \sum a_i$	1	2	1			1	2	1			
$(\sum a_i a_j a_k)^2$	1	1				1	1				
$\sum a_i a_j a_k a_l (\sum a_i)^2$		1				1	2	1			
$\sum a_i a_j a_k a_l \sum a_i a_m$		1				1	1				
$\sum a_i a_j a_k a_l \sum a_i$						1	1				
$\sum a_i a_j a_k a_l a_m a_n$						1					

	$\frac{10123 \dots (n-5)(n-3)(n-1)(n+1)}{10123 \dots (n-5)(n-3)(n-2)(n+1)}$ $\frac{10123 \dots (n-4)(n-1)(n+2)}{10123 \dots (n-5)(n-3)(n-2)(n+3)}$ $\frac{10123 \dots (n-4)(n-2)(n+2)}{10123 \dots (n-3)(n+3)}$ $\frac{10123 \dots (n-3)(n+3)}$ $\frac{10123 \dots (n-2)(n-9) \dots (n-1)}{10123 \dots (n-5)(n-3)(n-2)(n+3)}$ $\frac{10123 \dots (n-2)(n-1)(n+3)}{10123 \dots (n-2)(n+3)}$ $\frac{10123 \dots (n-2)(n-1)(n+3)}{10123 \dots (n-2)(n+3)}$														
$(\sum a_i)^7$	14	14	21	35	35	21	14	14	1	6	15	20	15	6	1
$\sum a_i a_j (\sum a_i)^5$	9	10	11	20	15	10	4	5	1	5	10	10	5	1	
$(\sum a_i a_j)^2 (\sum a_i)^3$	6	7	6	11	6	5	1	2	1	4	6	4	1		
$(\sum a_i a_j)^3 \sum a_i$	4	5	3	6	2	3		1	1	3	3	1			
$\sum a_i a_j a_k (\sum a_i)^4$	4	6	3	8	3	2			1	4	6	4	1		
$\sum a_i a_j a_k \sum a_i a_l (\sum a_i)^2$	3	4	2	4	1	1			1	3	3	1			
$\sum a_i a_j a_k (\sum a_i a_l)^2$	2	3	1	2		1			1	2	1				
$(\sum a_i a_j a_k)^2 \sum a_i$	2	2	1	1					1	2	1				
$\sum a_i a_j a_k a_l (\sum a_i)^3$	1	3		2					1	3	3	1			
$\sum a_i a_j a_k \sum a_i a_l \sum a_i$	1	2		1					1	2	1				
$\sum a_i a_j a_k a_l \sum a_i a_m a_n$	1	1							1	1					
$\sum a_i a_j a_k a_l a_m (\sum a_i)^2$		1							1	2	1				
$\sum a_i a_j a_k a_l a_m \sum a_i a_n$		1							1	1					
$\sum a_i a_j a_k a_l a_m a_n \sum a_i$									1	1					
$\sum a_i a_j a_k a_l a_m a_n a_p$									1						

4. If we denote the product

$$| 0123 \dots (n-1) | \left( \sum a_1 a_2 a_3 \dots a_{j_1} \right)^{i_2} \left( \sum a_1 a_2 a_3 \dots a_{j_2} \right)^{i_3} \dots \left( \sum a_1 a_2 a_3 \dots a_{j_h} \right)^{i_{h-1}}$$

by  $P$ , then the alternant

$| 0123 \dots r_1 s_1 \dots r_2 s_2 \dots r_g s_g \dots r_k s_k |$  —which we shall represent temporarily by  $A_1$ — in the product  $P \Sigma a_1 a_2 a_3 \dots a_{j_1}$ , (or  $P_1$ ), can arise from the following alternants of  $P$ :

$$| 0123 \dots r_1 (s_1 - 1) s_1 (s_1 + 1) \dots (s_1 + a_1 - 2) (s_1 + a_1) \dots r_g (s_g - 1) (s_g) \dots \\ \dots (s_g + r_1 - 2) (s_g + r_1) \dots r_k (s_k - 1) | ,$$

where  $\alpha_1, \beta_1, \dots, \gamma_1, \dots, \kappa_1 = 0, 1, 2, 3, \dots, j_1$ ,

and  $\alpha_1 + \beta_1 + \gamma_1 + \dots + \kappa_1 = j_1$ .

The alternant

$$| 0123 \dots (r_1 - j_h + 1) \dots r_1 (r_1 + 1) s_1 \dots r_2 s_2 \dots r_g s_g \dots r_k s_k | ,$$

which we shall also denote temporarily by  $A_2$  in the product  $P \Sigma a_1 a_2 a_3 \dots a_{j_1} + j_h$ , (or  $P_2$ ), will evidently arise from the same terms of  $P$ , but it will arise also from the following terms:

$$| 0123 \dots (r_1 - j_h + 1) \dots r_1 (r_1 + 1) s_1 (s_1 + 1) \dots (s_1 + a + a_1 - 2) (s_1 + a + a_1) \dots \\ \dots r_g s_g (s_g + 1) \dots (s_g + a + a_1 - 2) (s_g + a + a_1) \dots r_k (s_k - 1) | .$$

These terms could not exist, however, if  $j_1 \geq \frac{1}{2}n$ .

5. If we denote the coefficient of  $A_2$  by

$$C_{r_1}^{t j_h} \left\{ \begin{matrix} s_1 & s_2 & \dots & s_k \\ 0 & 0 & \dots & 0 \end{matrix} \right\}$$

where the  $j_h$  above the  $r_1$  denotes that  $r_1$  and the preceding  $(j_h - 1)$  numbers are increased by unity.

Then, if  $j_1 \geq \frac{1}{2}n$ , we will have

$$C_{r_1}^{t j_h} \left\{ \begin{matrix} s_1 & s_2 & \dots & s_k \\ 0 & 0 & \dots & 0 \end{matrix} \right\} = \sum_0^{j_h} \alpha \beta \dots \kappa C_{r_1}^{t - j_h} \left\{ \begin{matrix} s_1 & s_2 & \dots & s_k \\ \alpha & \beta & \dots & \kappa \end{matrix} \right\} ;$$

for, under these conditions the coefficient of  $A_1$  in the product  $P_1$  is the same as the coefficient of  $A_2$  in  $P_2$ .



[illegible]



9. We wish to prove the formula

$$C_{s_1}^{i0} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\} = \frac{i(i-1) \dots (i-s_1+2)}{\underline{s_1-1}} \cdot \frac{(s_1-2)(s_2-3) \dots (s_1-s_3+r_3-2)}{\underline{s_3-r_3-1}}.$$

In the reduction formula of article 8, let  $i$  receive the successive values 2, 3, 4, 5..... $i$ , and we have the following result :

$$C_{s_1}^2 0 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\} = C_{s_1}^1 1 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 1 \end{smallmatrix} \right\} + C_{s_1}^1 \left\{ \begin{smallmatrix} s_1 & s_2 \\ 1 & 0 \end{smallmatrix} \right\} + C_{s_1}^1 0 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\}$$

$$C_{s_1}^3 0 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\} = C_{s_1}^2 1 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 1 \end{smallmatrix} \right\} + C_{s_1}^2 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 0 \end{smallmatrix} \right\} + C_{s_1}^2 0 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\}$$

$$C_{s_1}^4 0 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\} = C_{s_1}^3 1 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 1 \end{smallmatrix} \right\} + C_{s_1}^3 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 0 \end{smallmatrix} \right\} + C_{s_1}^3 0 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\}$$

$$C_{s_1}^5 0 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\} = C_{s_1}^4 1 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 1 \end{smallmatrix} \right\} + C_{s_1}^4 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 0 \end{smallmatrix} \right\} + C_{s_1}^4 0 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\}$$

.....

$$C_{s_1}^{i-1} 0 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\} = C_{s_1}^{i-2} 1 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 1 \end{smallmatrix} \right\} + C_{s_1}^{i-2} \left\{ \begin{smallmatrix} s_2 & s_2 \\ 1 & 0 \end{smallmatrix} \right\} + C_{s_1}^{i-2} 0 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\}$$

$$C_{s_1}^{i0} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\} = C_{s_1}^{i-1} 1 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 1 \end{smallmatrix} \right\} + C_{s_1}^{i-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 0 \end{smallmatrix} \right\} + C_{s_1}^{i-1} 0 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\}$$

---


$$C_{s_1}^{i0} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\} = \sum_1^{i-1} j C_{s_1}^j 1 \left\{ \begin{smallmatrix} s_2 & s_2 \\ 0 & 1 \end{smallmatrix} \right\} + \sum_1^{i-1} j C_{s_1}^j 1 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 0 \end{smallmatrix} \right\}$$

10. Let the reduction formula, designated  $R$ ,

$$C_{s_1}^{i0} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\} = \sum_1^{i-1} j C_{s_1}^j 1 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 1 \end{smallmatrix} \right\} + \sum_1^{i-1} j C_{s_1}^j 1 \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 0 \end{smallmatrix} \right\}$$

operate upon itself and the successive results :

$$C^{i-0}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 0 \end{smallmatrix} \right\} = C^{i-1}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 1 \end{smallmatrix} \right\} + C^{i-1}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 0 \end{smallmatrix} \right\}$$

$$C^{i-2}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 1 \end{smallmatrix} \right\} + C^{i-2}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 0 \end{smallmatrix} \right\}$$

$$C^{i-3}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 1 \end{smallmatrix} \right\} + C^{i-3}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 0 \end{smallmatrix} \right\}$$

.....

$$C^{s_1-2}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 1 \end{smallmatrix} \right\} + C^{s_1-2}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 0 \end{smallmatrix} \right\}$$

Observe that the terms of  $R$  beyond the limit  $s_1 - 2$  are zero.

$$= 1 \left( C^{i-2}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 2 & 0 \end{smallmatrix} \right\} + 2 C^{i-2}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 1 \end{smallmatrix} \right\} + C^{i-2}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 2 \end{smallmatrix} \right\} \right)$$

$$2 \left( C^{i-3}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 2 & 0 \end{smallmatrix} \right\} + 2 C^{i-3}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 1 \end{smallmatrix} \right\} + C^{i-3}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 2 \end{smallmatrix} \right\} \right)$$

$$(a) \quad 3 \left( C^{i-4}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 2 & 0 \end{smallmatrix} \right\} + 2 C^{i-4}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 1 \end{smallmatrix} \right\} + C^{i-4}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 2 \end{smallmatrix} \right\} \right)$$

.....

$$(i-2) C^{s_1-3}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 2 & 0 \end{smallmatrix} \right\} + 2 C^{s_1-3}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 1 \end{smallmatrix} \right\} + C^{s_1-3}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 2 \end{smallmatrix} \right\}$$

$$= 1 \left( C^{i-3}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 3 & 0 \end{smallmatrix} \right\} + 3 C^{i-3}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 2 & 1 \end{smallmatrix} \right\} + 3 C^{i-3}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 2 \end{smallmatrix} \right\} + C^{i-3}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 3 \end{smallmatrix} \right\} \right)$$

$$3 \left( C^{i-4}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 3 & 0 \end{smallmatrix} \right\} + 3 C^{i-4}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 2 & 1 \end{smallmatrix} \right\} + 3 C^{i-4}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 2 \end{smallmatrix} \right\} + C^{i-4}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 3 \end{smallmatrix} \right\} \right)$$

$$(b) 6 \left( C^{i-5}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 3 & 0 \end{smallmatrix} \right\} + 3 C^{i-5}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 2 & 1 \end{smallmatrix} \right\} + 3 C^{i-5}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 2 \end{smallmatrix} \right\} + C^{i-5}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 3 \end{smallmatrix} \right\} \right)$$

$$10 \left( C^{i-6}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 3 & 0 \end{smallmatrix} \right\} + 3 C^{i-6}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 2 & 1 \end{smallmatrix} \right\} + 3 C^{i-6}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 1 & 2 \end{smallmatrix} \right\} + C^{i-6}_{s_1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 0 & 3 \end{smallmatrix} \right\} \right)$$

.....

$$\begin{aligned}
& \frac{(i-2)(i-3)}{\underline{2}} \left( C^{s_1-43} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & 0 \end{smallmatrix} \right\} + 3 C^{s_1-43} \left\{ \begin{smallmatrix} s_2 & s_3 \\ 2 & 1 \end{smallmatrix} \right\} \right. \\
& \qquad \qquad \qquad \left. + 3 C^{s_1-43} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & 1 \end{smallmatrix} \right\} + C^{s_1-43} \left\{ \begin{smallmatrix} s_2 & c_3 \\ 0 & 3 \end{smallmatrix} \right\} \right) \\
& \dots \qquad \dots \qquad \dots \qquad \dots \qquad \dots \qquad \dots \qquad \dots \qquad \dots \\
& \dots \qquad \dots \qquad \dots \qquad \dots \qquad \dots \qquad \dots \qquad \dots \qquad \dots \\
& = 1 \left( C^{i-s_1+2s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -2 \ 0 \end{smallmatrix} \right\} + (s_1-2) C^{i-s_1+2s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -3 \ 1 \end{smallmatrix} \right\} \right. \\
& \qquad \qquad \qquad \left. + \frac{(s_1-2)(s_1-3)}{\underline{2}} C^{i-s_1+2s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -4 \ 2 \end{smallmatrix} \right\} + \dots \right) \\
& (s_1-2) \left( C^{i-s_1+1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -2 \ 0 \end{smallmatrix} \right\} + (s_1-2) C^{i-s_1+1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -3 \ 1 \end{smallmatrix} \right\} \right. \\
& \qquad \qquad \qquad \left. + \frac{(s_1-2)(s_1-3)}{\underline{2}} C^{i-s_1+1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -4 \ 2 \end{smallmatrix} \right\} + \dots \right) \\
& (c) \frac{(s_1-2)(s_1-3)}{\underline{2}} \left( C^{i-s_1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -2 \ 0 \end{smallmatrix} \right\} + (s_1-2) C^{i-s_1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -3 \ 1 \end{smallmatrix} \right\} \right. \\
& \qquad \qquad \qquad \left. + \frac{(s_1-2)(s_1-3)}{\underline{2}} C^{i-s_1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -4 \ 2 \end{smallmatrix} \right\} + \dots \right) \\
& \frac{(s_1-2)(s_1-3)(s_1-4)}{\underline{3}} \left( C^{i-s_2-1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -2 \ 0 \end{smallmatrix} \right\} \right. \\
& \qquad \qquad \qquad \left. + (s_1-2) C^{i-s_2-1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -3 \ 1 \end{smallmatrix} \right\} \right. \\
& \qquad \qquad \qquad \left. + \frac{(s_1-2)(s_1-3)}{\underline{2}} C^{i-s_2-1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -4 \ 2 \end{smallmatrix} \right\} + \dots \right) \\
& \dots \qquad \dots \qquad \dots \qquad \dots \qquad \dots \qquad \dots \qquad \dots \qquad \dots \\
& \frac{(i-2)(i-3)\dots(i-s_1+2)}{\underline{s_1-3}} \left( C^{1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -2 \ 0 \end{smallmatrix} \right\} + (s_1-2) C^{1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1 & -3 \ 1 \end{smallmatrix} \right\} \right.
\end{aligned}$$

$$+ \frac{(s_1-2)(s_1-3)}{\underline{2}} C^{1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1-4 & 2 \end{smallmatrix} \right\} + \dots + \dots$$

$$+ \frac{(s_1-2)(s_1-3)\dots(s_1-k-1)}{\underline{k}} C^{1s_1-1} \left\{ \begin{smallmatrix} s_2 & s_3 \\ s_1-k-2 & k \end{smallmatrix} \right\} + \dots).$$

In (c) let  $i=i$ ,  $s_1=s_1$ , and  $s_3=r_3+1$ . Then

$$C^{i0} \left\{ \begin{smallmatrix} s_2 & (r_3+1) \\ 0 & 0 \end{smallmatrix} \right\} = 1 C^{i-s_1+2s_1-1} \left\{ \begin{smallmatrix} s_2 & (r_3+1) \\ s_1-2 & 0 \end{smallmatrix} \right\} +$$

$$(s_1-2) C^{i-s_1+1s_1-1} \left\{ \begin{smallmatrix} s_2 & (r_3+1) \\ s_1 & s_1-2 \end{smallmatrix} \right\} + \frac{(s_1-1)(s_1-2)}{\underline{2}} C^{i-s_1s_1-1} \left\{ \begin{smallmatrix} s_2 & (r_3+1) \\ s_1 & s_2-1 \end{smallmatrix} \right\}$$

$$+ \dots + \frac{(i-2)(i-3)(i-4)\dots(i-s_1+2)}{\underline{s_1-3}} C^{1s_1-1} \left\{ \begin{smallmatrix} s_2 & (r_3+1) \\ s_1 & s_1-1 \end{smallmatrix} \right\}.$$

It is evident from the table that the  $O$ 's have values, respectively, from  $i-s_1+2$  down to 1. Wherefore

$$C^{i0} \left\{ \begin{smallmatrix} s_2 & (r_3+1) \\ 0 & 0 \end{smallmatrix} \right\} = 1 \cdot (i-s_1+2) + (s_1-2)(i-s_1+1) + \frac{(s_1-1)(s_1-2)}{\underline{2}} (i-s_1)$$

$$+ \frac{s_1(s_1-1)(s_1-2)}{\underline{3}} (i-s_1-1) + \dots + \frac{(i-2)(i-3)(i-4)\dots(i-s_1+2)}{\underline{s_1-3}} (i-(i-1))$$

$$= i \left[ 1 + (s_1-2) + \frac{(s_1-1)(s_1-2)}{\underline{2}} + \frac{s_1(s_1-1)(s_1-2)}{\underline{3}} + \dots + \frac{(i-2)(i-3)\dots(i-s_1+2)}{\underline{s_1-3}} \right] - (s_1-2) \left[ 1 + (s_1-1) \right.$$

$$+ \frac{(s_1-1)s_1}{\underline{2}} + \frac{(s_1-1)s_1(s_1+1)}{\underline{3}} + \dots + \frac{(i-1)(i-2)(i-3)\dots(i-s_1+2)}{\underline{s_1+2}} \left. \right].$$

Summing the two series in the brackets, we have

$$C^{i0} \left\{ \begin{smallmatrix} s_2 & (r_3+1) \\ 0 & 0 \end{smallmatrix} \right\} = \frac{i(i-1)(i-2)\dots(i-s_1+2)}{\underline{s_1-2}} - (s_1-2) \frac{i(i-1)\dots(i-s_1+2)}{\underline{s_1-1}}$$

$$\begin{aligned}
&= \frac{i(i-1)(i-2)(i-3)\dots(i-s_1+2)}{|s_1-1|} (s_1-1-(s_1-2)) \\
&= \frac{i(i-1)(i-2)(i-3)\dots(i-s_1+2)}{|s_1-1|} .1.
\end{aligned}$$

In a similar manner, it can be proven that

$$\begin{aligned}
C_{s_1}^{i \ 0} \left\{ \begin{matrix} s_2 & r_3+2 \\ 0 & 0 \end{matrix} \right\} &= \frac{i(i-1)(i-2)(i-3)\dots(i-s_1+2)}{|s_1-1|} (s_1-2) \\
C_{s_1}^{i \ 0} \left\{ \begin{matrix} s_2 & r_3+3 \\ 0 & 0 \end{matrix} \right\} &= \frac{i(i-1)(i-2)(i-3)\dots(i-s_1+2)}{|s_1-1|} \frac{(s_1-2)}{2} \frac{(s_1-3)}{2},
\end{aligned}$$

or, in general,

$$\begin{aligned}
C_{s_1}^{i \ 0} \left\{ \begin{matrix} s_2 & s_3 \\ 0 & 0 \end{matrix} \right\} &= \frac{i(i-1)(i-2)(i-3)\dots(i-s_1+2)}{|s_1-1|} \\
&\times \frac{(s_1-2)(s_1-3)\dots(s_1-s_3+r_3-2)}{|s_3-r_3-1|}.
\end{aligned}$$

SYRACUSE UNIVERSITY, *December 18, 1902.*

---

## AN ACCOUNT OF PROFESSOR RUNKLE'S MATHEMATICAL MONTHLY.

---

By PROFESSOR SIMON NEWCOMB.

---

I first made Mr. Runkle's acquaintance in the winter of 1857, when he was senior assistant in the Nautical Almanac Office, then at Cambridge, Mass. His intelligence, intellectual activity, and lively interest in matters and things generally, not excluding things political, made him a very interesting character.

It was early in 1858 that he announced to me and some others in the office his intention of starting a mathematical journal. His first step was to secure the necessary support. It may be feared that few in our day have an adequate conception of the backward condition of mathematical study in our country at that time. A curious illustration is offered by Davies' well-known dictionary of mathematics, in the preface of which it was announced that it contained defin-